



VASN Polyclonal Antibody

Catalog No	YP-Ab-06905
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	VASN SLITL2 UNQ314/PRO357/PRO1282
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	VASN Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	VASN;VASN SLITL2 UNQ314/PRO357/PRO1282;Vasorin (Protein slit-like 2)
Observed Band	74kD
Calculated Molecular Weight	74kD
Cell Pathway	Membrane ; Single-pass type I membrane protein . Secreted .
Tissue Specificity	Expressed at highest levels in aorta, at intermediate levels in kidney and placenta and at lowest levels in brain, heart, liver, lung and skeletal muscle. Within the aorta, the strongest expression is found in the tunica media of the proximal ascending aorta, the descending thoracic aorta, the abdominal aorta and the coronary arteries. Within the kidney, expression is found in the interstitial cells.
Function	function:May act as an inhibitor of TGF-beta signaling.,PTM:N-glycosylated.,similarity:Contains 1 EGF-like domain.,similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 11 LRR (leucine-rich) repeats.,subunit:Interacts with TGFB1, TGFB2 and TGFB3.,tissue specificity:Expressed at highest levels in aorta, at intermediate levels in kidney and placenta and at lowest levels in brain, heart, liver, lung and skeletal muscle. Within the aorta, the strongest expression is found in the tunica media of the proximal ascending aorta, the descending thoracic aorta, the abdominal aorta and the coronary arteries. Within the kidney,

expression is found in the interstitial cells.,

Background

function:May act as an inhibitor of TGF-beta signaling.,PTM:N-glycosylated.,
similarity:Contains 1 EGF-like domain.,similarity:Contains 1 fibronectin type-III
domain.,similarity:Contains 11 LRR (leucine-rich) repeats.,subunit:Interacts with
TGFB1, TGFB2 and TGFB3.,tissue specificity:Expressed at highest levels in
aorta, at intermediate levels in kidney and placenta and at lowest levels in brain,
heart, liver, lung and skeletal muscle. Within the aorta, the strongest expression
is found in the tunica media of the proximal ascending aorta, the descending
thoracic aorta, the abdominal aorta and the coronary arteries. Within the kidney,
expression is found in the interstitial cells.,

**matters needing
attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For
more information, please consult technical personnel.

Products Images