



ROBO3 Polyclonal Antibody

Catalog No	YP-Ab-07044
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	ROBO3
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	ROBO3 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ROBO3;HGPPS;HGPS;RBIG1;RIG1;Roundabout homolog 3 (Roundabout-like protein 3)
Observed Band	148 kDa
Calculated Molecular Weight	148 kDa
Cell Pathway	Membrane ; Single-pass type I membrane protein .
Tissue Specificity	Brain,
Function	disease:Defects in ROBO3 are a cause of familial horizontal gaze palsy with progressive scoliosis (HGPPS) [MIM:607313]. Patients show a medulla where motor and sensory projections appear uncrossed.,function:Thought to be involved during neural development in axonal navigation at the ventral midline of the neural tube. In spinal chord development plays a role in guiding commissural axons probably by preventing premature sensitivity to Slit proteins thus inhibiting Slit signaling through ROBO1 (By similarity). Required for hindbrain axon midline crossing.,similarity:Belongs to the immunoglobulin superfamily. ROBO family.,similarity:Contains 3 fibronectin type-III domains.,similarity:Contains 5 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Probably interacts with SLIT2.,

Background

This gene is a member of the Roundabout (ROBO) gene family that controls neurite outgrowth, growth cone guidance, and axon fasciculation. ROBO proteins are a subfamily of the immunoglobulin transmembrane receptor superfamily. SLIT proteins 1-3, a family of secreted chemorepellants, are ligands for ROBO proteins and SLIT/ROBO interactions regulate myogenesis, leukocyte migration, kidney morphogenesis, angiogenesis, and vasculogenesis in addition to neurogenesis. This gene, ROBO3, has a putative extracellular domain with five immunoglobulin (Ig)-like loops and three fibronectin (Fn) type III motifs, a transmembrane segment, and a cytoplasmic tail with three conserved signaling motifs: CC0, CC2, and CC3 (CC for conserved cytoplasmic). Unlike other ROBO family members, ROBO3 lacks motif CC1. The ROBO3 gene regulates axonal navigation at the ventral midline of the neural tube. In mouse, loss of R

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images