



LGMN Polyclonal Antibody

Catalog No	YP-Ab-07239
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	LGMN PRSC1
Immunogen	Synthesized peptide derived from human protein . at AA range: 60-140
Specificity	LGMN Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	1F2B12;EC:3.4.22.34;legumain;PRSC1;LGMN;LGMN PRSC1;Legumain (EC 3.4.22.34) (Asparaginyl endopeptidase) (Protease;cysteine 1)
Observed Band	47kD
Calculated Molecular Weight	47kD
Cell Pathway	Lysosome .
Tissue Specificity	Ubiquitous. Particularly abundant in kidney, heart and placenta.
Function	catalytic activity:Hydrolysis of proteins and small molecule substrates at -Asn- -Xaa- bonds.,function:Has a strict specificity for hydrolysis of asparaginyl bonds. Can also cleave aspartyl bonds slowly, especially under acidic conditions. May be involved in the processing of proteins for MHC class II antigen presentation in the lysosomal/endosomal system.,PTM:Glycosylated.,similarity:Belongs to the peptidase C13 family.,tissue specificity:Ubiquitous. Particularly abundant in kidney, heart and placenta.,
Background	This gene encodes a cysteine protease that has a strict specificity for hydrolysis of asparaginyl bonds. This enzyme may be involved in the processing of bacterial peptides and endogenous proteins for MHC class II presentation in the lysosomal/endosomal systems. Enzyme activation is triggered by acidic pH and



appears to be autocatalytic. Protein expression occurs after monocytes differentiate into dendritic cells. A fully mature, active enzyme is produced following lipopolysaccharide expression in mature dendritic cells. Overexpression of this gene may be associated with the majority of solid tumor types. This gene has a pseudogene on chromosome 13. Several alternatively spliced transcript variants have been described, but the biological validity of only two has been determined. These two variants encode the same isoform. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night