



MYH9 Polyclonal Antibody

Catalog No	YP-Ab-07818
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	MYH9
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	MYH9 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Myosin 9;Myosin heavy chain 9;Myosin heavy chain;non-muscle IIa;Myosin-9; NMMHC II-a;MYH9;Myosin-9 (Cellular myosin heavy chain;type A) (Myosin heavy chain 9) (Myosin heavy chain;non-muscle IIa) (Non-muscle myosin heavy chain A) (NMMHC-A) (Non-muscle myosin heavy chain IIa) (NMMHC II-a)
Observed Band	224 kDa
Calculated Molecular Weight	227 kDa
Cell Pathway	Cytoplasm, cytoskeleton . Cytoplasm, cell cortex . Cytoplasmic vesicle, secretory vesicle, Cortical granule . Colocalizes with actin filaments at lamellipodia margins and at the leading edge of migrating cells (PubMed:20052411). In retinal pigment epithelial cells, predominantly localized to stress fiber-like structures with some localization to cytoplasmic puncta (PubMed:27331610). .
Tissue Specificity	In the kidney, expressed in the glomeruli. Also expressed in leukocytes.
Function	disease:Defects in MYH9 are the cause of Alport syndrome with macrothrombocytopenia (APSM) [MIM:153650]. APSM is an autosomal dominant disorder characterized by the association of ocular lesions, sensorineural hearing loss and nephritis (Alport syndrome) with platelet defects., disease:Defects in MYH9 are the cause of Epstein syndrome (EPS)



[MIM:153650]. EPS is an autosomal dominant disorder characterized by the association of macrothrombocytopathy, sensorineural hearing loss and nephritis., disease:Defects in MYH9 are the cause of Fechtner syndrome (FTNS) [MIM:153640]. FTNS is an autosomal dominant macrothrombocytopenia characterized by thrombocytopenia, giant platelets and leukocyte inclusions that are small and poorly organized. Additionally, FTNS is distinguished by Alport-like clinical features of sensorineural deafness, cataracts and nephritis., disease:Defects in MYH9 are the cause o

Background

This gene encodes a conventional non-muscle myosin; this protein should not be confused with the unconventional myosin-9a or 9b (MYO9A or MYO9B). The encoded protein is a myosin IIA heavy chain that contains an IQ domain and a myosin head-like domain which is involved in several important functions, including cytokinesis, cell motility and maintenance of cell shape. Defects in this gene have been associated with non-syndromic sensorineural deafness autosomal dominant type 17, Epstein syndrome, Alport syndrome with macrothrombocytopenia, Sebastian syndrome, Fechtner syndrome and macrothrombocytopenia with progressive sensorineural deafness. [provided by RefSeq, Dec 2011],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images