



YTHD2 Polyclonal Antibody

Catalog No	YP-Ab-07868
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	YTHDF2 HGRG8
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	YTHD2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	YTH domain family protein 2 (CLL-associated antigen KW-14) (High-glucose-regulated protein 8) (Renal carcinoma antigen NY-REN-2);YTHD2;YTHDF2 HGRG8
Observed Band	63kD
Calculated Molecular Weight	63kD
Cell Pathway	Cytoplasm, cytosol . Cytoplasm, P-body . Cytoplasm, Stress granule . Nucleus . Localizes to the cytosol and relocates to the nucleus following heat shock stress (PubMed:26458103). Can partition into different structures: into P-bodies in unstressed cells, and into stress granules during stress (PubMed:31292544). .
Tissue Specificity	Highly expressed in induced pluripotent stem cells (iPSCs) and down-regulated during neural differentiation.
Function	similarity:Contains 1 YTH domain.,
Background	This gene encodes a member of the YTH (YT521-B homology) superfamily containing YTH domain. The YTH domain is typical for the eukaryotes and is particularly abundant in plants. The YTH domain is usually located in the middle of the protein sequence and may function in binding to RNA. In addition to a YTH domain, this protein has a proline rich region which may be involved in

signal transduction. An Alu-rich domain has been identified in one of the introns of this gene, which is thought to be associated with human longevity. In addition, reciprocal translocations between this gene and the Runx1 (AML1) gene on chromosome 21 has been observed in patients with acute myeloid leukemia. This gene was initially mapped to chromosome 14, which was later turned out to be a pseudogene. Alternatively spliced transcript variants encoding different isoforms have been identified in this gene

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images