



TRI41 rabbit pAb


Catalog No	YP-Ab-08329
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	TRIM41 RINCK
Immunogen	Synthesized peptide derived from human TRI41 AA range: 464-514
Specificity	This antibody detects endogenous levels of TRI41 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	E3 ubiquitin-protein ligase TRIM41;MGC1127;MGC31991;RINCK;RING finger protein that interacts with C kinase;RING finger-interacting protein with C kinase; TRI41_HUMAN;Trim41;Tripartite motif containing 41;Tripartite motif-containing protein 41;TRI41;TRIM41 RINCK
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Cytoplasm . Nucleus .
Tissue Specificity	Expressed in multiple tissues with the highest levels in heart and skeletal muscle.
Function	similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B box-type zinc finger.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 RING-type zinc finger.,
Background	This gene encodes a member of the tripartite motif (TRIM) family. The TRIM family is characterized by a signature motif composed of a RING finger, one or more B-box domains, and a coiled-coil region. This encoded protein may play a role in protein kinase C signaling. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products ImagesWestern blot analysis of lysates from MCF-7 cells,
primary antibody was diluted at 1:1000, 4° over night