



GSX1 rabbit pAb

Catalog No	YP-Ab-08510
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	GSX1 GSH1
Immunogen	Synthesized peptide derived from human GSX1 AA range: 131-181
Specificity	This antibody detects endogenous levels of GSX1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ABC 29;ABC29;ABCC 1;ABCC;Abcc1;ATP binding cassette sub family C (CFTR/MRP) member 1;ATP binding cassette sub-family C member 1;ATP binding cassette subfamily C member 1;ATP binding cassette transporter variant ABCC1delta ex13;ATP binding cassette transporter variant ABCC1delta ex13&14;ATP binding cassette transporter variant ABCC1delta ex25;ATP binding cassette transporter variant ABCC1delta ex25&26;ATP binding cassette; sub-family C (CFTR/MRP);member 1;ATP-binding cassette sub-family C member 1;DKFZp686N4233;DKFZp781G125;GS X;GSX;Leukotriene C(4) transporter;LTC4 transporter;MRP 1;MRP;MRP1;MRP1_HUMAN;Multidrug resistance associated protein 1;Multidrug resistance protein;Multidrug resistance-associated protein 1;Multiple drug resistance associated protein;Multiple drug resistance protein 1;GSX1;GSX1 GSH1
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Nucleus .
Tissue Specificity	




Function function: Probable transcription factor that binds to the DNA sequence 5'-GC[TA][AC]ATTA[GA]-3'. Activates the transcription of the GHRH gene. Plays an important role in pituitary development., similarity: Belongs to the Antp homeobox family., similarity: Contains 1 homeobox DNA-binding domain.,

Background

matters needing attention Avoid repeated freezing and thawing!

Usage suggestions This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night