



1A68 rabbit pAb

Catalog No	YP-Ab-08949
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	HLA-A HLAA
Immunogen	Synthesized peptide derived from human 1A68 AA range: 1-51
Specificity	This antibody detects endogenous levels of 1A68 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	1A68;HLA-A HLAA
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Golgi membrane,endoplasmic reticulum,Golgi apparatus,Golgi medial cisterna, plasma membrane,integral component of plasma membrane,cell surface,ER to Golgi transport vesicle membrane,membrane,integral component of membrane,
Tissue Specificity	
Function	function:Involved in the presentation of foreign antigens to the immune system., polymorphism:The following alleles of A-1 are known: A*0101, A*0102, A*0103, A*0106 and A*0107. The sequence shown is that of A*0101.,polymorphism:The following alleles of A-11 are known: A*1101 (A-11E), A*1102 (A-11K), A*1103, A*1104, A*1105 and A*1107. The sequence shown is that of A*1101., polymorphism:The following alleles of A-2 are known: A*0201, A*0202, A*0203, A*0204, A*0205, A*0206 (A2.4A), A*0207, A*0208, A*0209, A*0210, A*0211 (A2.5), A*0212, A*0213 (A*02SLU), A*0216, A*0217, A*0218 (A2K), A*0219, A*0220, A*0221, A*0231, A*0234 (A*AAT), A*0235, A*0236 and A*0237. The sequence shown is that of A*0201.,polymorphism:The following alleles of A-23 are known: A*2301, A*2302, A*2303, A*2304 and A*2305. The sequence shown



is that of A*2301.,polymorphism:The following alleles of A-24 are known: A*2401, A*240

Background

HLA-A belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domains, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is routinely done for bone marrow and kidney trans


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night