



CP4AB rabbit pAb

Catalog No	YP-Ab-08985
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CYP4A11 CYP4A2
Immunogen	Synthesized peptide derived from human CP4AB AA range: 389-439
Specificity	This antibody detects endogenous levels of CP4AB at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	2-HETE synthase;2-hydroxyeicosatetraenoic acid synthase;CP4AB_HUMAN; CYP4A;Cyp4a1;CYP4A11;Cyp4a14;Cyp4a3;CYP4A7;CYP4A11;CYP4A11; Cytochrome P-45HK-omega;Cytochrome P45 4A1;Cytochrome P45 4A11; Cytochrome P45 4A14;Cytochrome P45 4A2;Cytochrome P45 4A3;Cytochrome P45 4A7;Cytochrome P45HL-omega;Fatty acid omega-hydroxylase;Lauric acid omega-hydroxylase;CP4AB;CYP4A11 CYP4A2
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Endoplasmic reticulum membrane ; Peripheral membrane protein . Microsome membrane ; Peripheral membrane protein .
Tissue Specificity	Expressed in liver (PubMed:7679927). Expressed in S2 and S3 segments of proximal tubules in cortex and outer medulla of kidney (PubMed:7679927, PubMed:10660572).
Function	catalytic activity:Octane + reduced rubredoxin + O(2) = 1-octanol + oxidized rubredoxin + H(2)O.,cofactor:Heme group.,function:Catalyzes the omega- and (omega-1)-hydroxylation of various fatty acids such as laurate, myristate and palmitate. Has little activity toward prostaglandins A1 and E1. Oxidizes arachidonic acid to 20-hydroxyeicosatetraenoic acid (20-HETE).,online



information:CYP4A11 alleles,polymorphism:CYP4A11v seems to be a rare allelic variant of CYP4A11, it seems to be unstable and not to metabolize lauric acid.,similarity:Belongs to the cytochrome P450 family.,tissue specificity:Kidney and liver.,

Background

cytochrome P450 family 4 subfamily A member 11(CYP4A11) Homo sapiens This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and hydroxylates medium-chain fatty acids such as laurate and myristate. Multiple transcript variants have been found for this gene. [provided by RefSeq, Jan 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night