



ASGL1 rabbit pAb

Catalog No	YP-Ab-09177
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	ASRGL1 ALP CRASH
Immunogen	Synthesized peptide derived from human ASGL1 AA range: 57-107
Specificity	This antibody detects endogenous levels of ASGL1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ALP; ALP1; ASGL1; Asparaginase like 1; Asparaginase like 1 protein; Asparaginase like protein 1; ASRGL 1; beta-aspartyl-peptidase; CRASH; FLJ22316; isoaspartyl dipeptidase; isoaspartyl peptidase/L-asparaginase; L asparaginase; L asparagine amidohydrolase; ASRGL1 ALP CRASH
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Cytoplasm . Midpiece of sperm tail.
Tissue Specificity	Expressed in brain, kidney, testis and tissues of the gastrointestinal tract. Present in sperm (at protein level). Over-expressed in uterine, mammary, prostatic and ovarian carcinoma.
Function	catalytic activity: L-asparagine + H ₂ O = L-aspartate + NH ₃ ., function: Acts in asparagine catabolism. May be involved in astroglial production of L-aspartate, which can act as an excitatory neurotransmitter in some brain regions., induction: By 5-alpha-di-hydrotestosterone and progesterone., similarity: Belongs to the Ntn-hydrolase family., subcellular location: Midpiece of sperm tail., tissue specificity: Expressed in brain, kidney, testis and tissues of the gastrointestinal tract. Present in sperm (at protein level). Over-expressed in uterine, mammary, prostatic and ovarian carcinoma.,

Background


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night