



NACA2 rabbit pAb

Catalog No	YP-Ab-09730
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	NACA2 NACAL
Immunogen	Synthesized peptide derived from human NACA2 AA range: 58-108
Specificity	This antibody detects endogenous levels of human NACA2
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Alpha NAC;Alpha-NAC;Hom s 2.2;HSD48;MGC117224;NAC-alpha;naca;NACA/1.9.2;NACA_HUMAN;NACA1;Nascent polypeptide associated complex alpha polypeptide;Nascent polypeptide associated complex alpha subunit;Nascent polypeptide associated complex subunit alpha;Nascent polypeptide-associated complex subunit alpha;Alpha NAC like;ANAC;Hom s 2.1;MGC71999;NAC alpha like;NACA2;NACAL;nascent polypeptide associated complex alpha polypeptide like;Nascent polypeptide associated complex subunit alpha 2;NACA2 NACAL;Nascent polypeptide-associated complex subunit alpha-2 (Alpha-NAC-like) (Hom s 2.01) (Nascent polypeptide-associated complex subunit alpha-like) (NAC-alpha-like)
Observed Band	
Calculated Molecular Weight	24KD; 23kD
Cell Pathway	Cytoplasm . Nucleus .
Tissue Specificity	Expressed specifically in testis and skeletal muscle.
Function	
Background	

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images

