



COX1 Monoclonal Antibody

Catalog No	YP-mAb-02383
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PTGS1
Immunogen	Synthetic Peptide of COX1 AA range: 483-533
Specificity	The antibody detects endogenous COX1 proteins.
Formulation	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PGH1;PTGS1;PGHS-1;EC:1.14.99.1;Cyclooxygenase-1;COX1;Prostaglandin G/H synthase 1;COX-1;Prostaglandin H2 synthase 1;PGH synthase 1;PHS 1;Prostaglandin-endoperoxide synthase 1
Observed Band	70kD
Calculated Molecular Weight	70kD
Cell Pathway	Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein.
Tissue Specificity	Brain,Lung fibroblast,Platelet,
Function	catalytic activity:Arachidonate + AH(2) + 2 O(2) = prostaglandin H(2) + A + H(2) O.,cofactor:Binds 1 heme B (iron-protoporphyrin IX) group per subunit.,function:May play an important role in regulating or promoting cell proliferation in some normal and neoplastically transformed cells.,miscellaneous:This enzyme acts both as a dioxygenase and as a peroxidase.,miscellaneous:This enzyme is the target of nonsteroidal anti-inflammatory drugs such as aspirin.,pathway:Lipid metabolism; prostaglandin biosynthesis.,similarity:Belongs to the prostaglandin G/H synthase family.,similarity:Contains 1 EGF-like domain.,subunit:Homodimer.,



Background

This is one of two genes encoding similar enzymes that catalyze the conversion of arachinodate to prostaglandin. The encoded protein regulates angiogenesis in endothelial cells, and is inhibited by nonsteroidal anti-inflammatory drugs such as aspirin. Based on its ability to function as both a cyclooxygenase and as a peroxidase, the encoded protein has been identified as a moonlighting protein. The protein may promote cell proliferation during tumor progression. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using COX1 Monoclonal Antibody