



PTH-rP Monoclonal Antibody

Catalog No	YP-mAb-10606
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PTH LH
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human PTH LH. AA range:101-150
Specificity	PTH-rP Monoclonal Antibody detects endogenous levels of PTH-rP
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PTH-rP;PTH LH;Parathyroid hormone-related protein (PTH-rP) (PTHrP) (Parathyroid hormone-like protein) (PLP) [Cleaved into: PTHrP[1-36];PTHrP[38-94];Osteostatin (PTHrP[107-139])]
Observed Band	25kD
Calculated Molecular Weight	25kD
Cell Pathway	Cytoplasm. Nucleus. Secreted.
Tissue Specificity	Ubiquitous. Also expressed in the mammary gland.
Function	alternative products:Additional isoforms seem to exist,disease:Produced by many tumors from patients with HHM (humoral hypercalcemia of malignancy)., function:Neuroendocrine peptide which is a critical regulator of cellular and organ growth, development, migration, differentiation and survival and of epithelial calcium ion transport. Regulates endochondral bone development and epithelial-mesenchymal interactions during the formation of the mammary glands and teeth.,function:Osteostatin is a potent inhibitor of osteoclastic bone resorption.,PTM:There are 3 principal secretory forms, called PTHrP[1-36], PTHrP[38-94], and osteostatin (PTHrP[107-139]) arising from endoproteolytic cleavage of the initial translation product. Each of these secretory forms is



believed to have one or more of its own receptors that mediates the normal paracrine, autocrine and endocrine actions.,similarity:Belongs

Background

parathyroid hormone like hormone(PTHrP) Homo sapiens The protein encoded by this gene is a member of the parathyroid hormone family. This hormone, via its receptor, PTHR1, regulates endochondral bone development and epithelial-mesenchymal interactions during the formation of the mammary glands and teeth. It is responsible for most cases of humoral hypercalcemia of malignancy, and mutations in this gene are associated with brachydactyly type E2 (BDE2). Alternatively spliced transcript variants have been found for this gene. There is also evidence for alternative translation initiation from non-AUG (CUG and GUG) start sites, downstream of the initiator AUG codon, resulting in nuclear forms of this hormone. [provided by RefSeq, Nov 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PTH-rP Monoclonal Antibody