



BLBP Monoclonal Antibody

Catalog No	YP-mAb-10653
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	FABP7 BLBP FABPB MRG
Immunogen	Synthetic peptide from human protein at AA range: 80-132
Specificity	The antibody detects endogenous BLBP
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	B FABP;BLBP;Brain lipid binding protein;FABP7;FABPB;Fatty acid binding protein 7;MRG;FABP7 BLBP FABPB MRG;Fatty acid-binding protein;brain (Brain lipid-binding protein) (BLBP) (Brain-type fatty acid-binding protein) (B-FABP) (Fatty acid-binding protein 7) (Mammary-derived growth inhibitor related); brain (Brain lipid-binding protein) (BLBP) (Brain-type fatty acid-binding protein) (B-FABP) (Fatty acid-binding protein 7) (Mammary-derived growth inhibitor related)
Observed Band	15kD
Calculated Molecular Weight	15kD
Cell Pathway	Cytoplasm.
Tissue Specificity	Expressed in brain and other neural tissues.
Function	domain:Forms a beta-barrel structure that accommodates the hydrophobic ligand in its interior.,function:B-FABP could be involved in the transport of a so far unknown hydrophobic ligand with potential morphogenic activity during CNS development. It is required for the establishment of the radial glial fiber system in developing brain, a system that is necessary for the migration of immature neurons to establish cortical layers.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,tissue specificity:Expressed in brain



and other neural tissues.,

Background

The gene encodes a small, highly conserved cytoplasmic protein that bind long-chain fatty acids and other hydrophobic ligands. The encoded protein is important in the establishment of the radial glial fiber in the developing brain. Alternative splicing and promoter usage results in multiple transcript variants encoding different isoforms. Pseudogenes of this gene are found on multiple chromosomes. [provided by RefSeq, Jan 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using BLBP Monoclonal Antibody