



PEK/PERK Monoclonal Antibody

Catalog No	YP-mAb-10796
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	EIF2AK3 PEK PERK
Immunogen	Synthesized peptide derived from human PEK/PERK Monoclonal
Specificity	This antibody detects endogenous levels of PEK/PERK.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EIF2AK3;PERK;EC:2.7.11.1;EIF2AK 3;Eukaryotic translation initiation factor 2-alpha kinase 3;EIF2AK3 PEK PERK;Eukaryotic translation initiation factor 2-alpha kinase 3 (EC 2.7.11.1) (PRKR-like endoplasmic reticulum kinase) (Pancreatic eIF2-alpha kinase) (HsPEK)
Observed Band	130kD
Calculated Molecular Weight	130kD
Cell Pathway	Endoplasmic reticulum membrane; Single-pass type I membrane protein.
Tissue Specificity	Ubiquitous. A high level expression is seen in secretory tissues.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.;disease:Defects in EIF2AK3 are the cause of Wolcott-Rallison syndrome (WRS) [MIM:226980]; also known as multiple epiphyseal dysplasia with early-onset diabetes mellitus. WRS is a rare autosomal recessive disorder, characterized by permanent neonatal or early infancy insulin-dependent diabetes and, at a later age, epiphyseal dysplasia, osteoporosis, growth retardation and other multisystem manifestations, such as hepatic and renal dysfunctions, mental retardation and cardiovascular abnormalities.;domain:The luminal domain senses perturbations in protein folding in the ER, probably through reversible interaction with HSPA5/BIP.;enzyme regulation:Perturbation in protein folding in the endoplasmic



reticulum (ER) promotes reversible dissociation from HSPA5/BIP and oligomerization, resulting in transautophosphorylation and kinase act

Background

The protein encoded by this gene phosphorylates the alpha subunit of eukaryotic translation-initiation factor 2, leading to its inactivation, and thus to a rapid reduction of translational initiation and repression of global protein synthesis. This protein is thought to modulate mitochondrial function. It is a type I membrane protein located in the endoplasmic reticulum (ER), where it is induced by ER stress caused by malfolded proteins. Mutations in this gene are associated with Wolcott-Rallison syndrome. [provided by RefSeq, Sep 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PEK/PERK Monoclonal Antibody