




TRI69 mouse mAb

Catalog No	YP-mAb-11381
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	TRIM69 RNF36 HSD-34 HSD34
Immunogen	Synthesized peptide derived from human TRI69 AA range: 202-252
Specificity	This antibody detects endogenous levels of TRI69 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, mouse,IgG
Purification	The antibody was affinity-purified from mouse serum by affinity-chromatography using specific immunogen.
Dilution	WB 1 :500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Observed Band	
Calculated Molecular Weight	55kD
Cell Pathway	Cytoplasm . Nucleus . Nucleus speckle .
Tissue Specificity	
Function	function:May play a role in apoptosis.,PTM:Phosphorylated. Phosphorylation is necessary for nuclear localization.,similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 RING-type zinc finger.,subunit:Interacts with PML.,
Background	This gene encodes a member of the RING-B-box-coiled-coil (RBCC) family and encodes a protein with an N-terminal RING finger motif, a PRY domain and a C-terminal SPRY domain. The mouse ortholog of this gene is specifically expressed in germ cells at the round spermatid stages during spermatogenesis and, when overexpressed, induces apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western blot analysis of lysates from AD293 cells, primary antibody was diluted at 1:1000, 4° over night