



PFD4 mouse mAb


Catalog No	YP-mAb-11652
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PFDN4 PFD4
Immunogen	Synthesized peptide derived from human PFD4 AA range: 65-115
Specificity	This antibody detects endogenous levels of PFD4 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	C1;PFD4;PFDN4;prefoldin 4;prefoldin subunit 4;Protein C 1;PFDN4 PFD4
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Nucleus . Cytoplasm . Mitochondrion .
Tissue Specificity	
Function	function: Binds specifically to cytosolic chaperonin (c-CPN) and transfers target proteins to it. Binds to nascent polypeptide chain and promotes folding in an environment in which there are many competing pathways for nonnative proteins., similarity: Belongs to the prefoldin subunit beta family., subunit: Heterohexamer of two PFD-alpha type and four PFD-beta type subunits.,
Background	This gene encodes a member of the prefoldin beta subunit family. The encoded protein is one of six subunits of prefoldin, a molecular chaperone complex that binds and stabilizes newly synthesized polypeptides, thereby allowing them to fold correctly. The complex, consisting of two alpha and four beta subunits, forms a double beta barrel assembly with six protruding coiled-coils. [provided by RefSeq, Jul 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using PFD4 mouse mAb