



SASH1 mouse mAb

Catalog No	YP-mAb-12061
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC
Gene Name	SASH1 KIAA0790 PEPE1
Immunogen	Synthesized peptide derived from human SASH1 AA range: 373-423
Specificity	This antibody detects endogenous levels of SASH1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, mouse,IgG
Purification	The antibody was affinity-purified from mouse serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DUH;DUH1;CAPOK;SH3D6A;dJ323M4.1;SASH1;SASH1 KIAA0790 PEPE1
Observed Band	137kD
Calculated Molecular Weight	
Cell Pathway	Cytoplasm .
Tissue Specificity	Expressed ubiquitously, with highest levels in lung, placenta, spleen and thymus. Down-regulated in the majority (74%) of breast tumors in comparison with corresponding normal breast epithelial tissues. Expressed in the epidermis, epidermal keratinocytes, dermal fibroblasts and melanocytes (PubMed:23333244, PubMed:26203640).
Function	function:May have a role in a signaling pathway. Could act as a tumor suppressor.,similarity:Contains 1 SH3 domain.,similarity:Contains 2 SAM (sterile alpha motif) domains.,tissue specificity:Expressed ubiquitously, with highest levels in lung, placenta, spleen and thymus. Down-regulated in the majority (74%) of breast tumors in comparison with corresponding normal breast epithelial tissues.,
Background	




matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).