



Cerebellin 4 Monoclonal Antibody

Catalog No	YP-mAb-12701
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CBLN4
Immunogen	The antiserum was produced against synthesized peptide derived from human CBLN4. AA range:141-190
Specificity	Cerebellin 4 Monoclonal Antibody detects endogenous levels of Cerebellin 4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CBLNL1;Cerebellin 4;cerebellin 4 precursor;Cerebellin like glycoprotein 1; Cerebellin-4;CBLN4;Cerebellin-like glycoprotein 1
Observed Band	27kD
Calculated Molecular Weight	27kD
Cell Pathway	Secreted . Cell junction, synapse . Detected at GABAergic synapses. .
Tissue Specificity	Brain,
Function	function:May be involved in synaptic functions in the CNS. Can enable ER export and secretion of CBLN3.,similarity:Contains 1 C1q domain., subunit:Homohexamer; disulfide-linked homotrimers. The trimers are assembled via the globular C1q domains. The trimers associate via N-terminal cysteine residues to form disulfide-linked hexamers. May interact with CBLN1, CBLN2 and CBLN3.,
Background	This gene encodes a member of a family of small secreted proteins containing C1Q domains. Members of this family are involved in regulation of neuroligin signalling during synapse development. The mouse homolog of the protein



encoded by this gene competes with netrin to bind to the deleted in colorectal cancer receptor. [provided by RefSeq, Aug 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Cerebellin 4 Monoclonal Antibody