



EAAT1 Monoclonal Antibody

Catalog No	YP-mAb-12718
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SLC1A3
Immunogen	The antiserum was produced against synthesized peptide derived from human EAAT1. AA range:492-541
Specificity	EAAT1 Monoclonal Antibody detects endogenous levels of EAAT1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EAAT1;GLAST;GLAST-1;SLC1A3;Sodium-dependent glutamate/aspartate transporter 1;Excitatory amino acid transporter 1;GLAST1;Solute carrier family 1 member 3
Observed Band	65kD
Calculated Molecular Weight	65kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Detected in brain (PubMed:8218410, PubMed:7521911, PubMed:8123008). Detected at very much lower levels in heart, lung, placenta and skeletal muscle (PubMed:7521911, PubMed:8123008). Highly expressed in cerebellum, but also found in frontal cortex, hippocampus and basal ganglia (PubMed:7521911).
Function	disease:Defects in SLC1A3 are the cause of episodic ataxia type 6 (EA6) [MIM:612656]. EA6 is characterized by episodic ataxia, seizures, migraine and alternating hemiplegia.,function:Transports L-glutamate and also L- and D-aspartate. Essential for terminating the postsynaptic action of glutamate by rapidly removing released glutamate from the synaptic cleft. Acts as a symport by cotransporting sodium.,PTM:Glycosylated.,similarity:Belongs to the sodium:dicarboxylate (SDF) symporter (TC 2.A.23) family.,tissue



specificity: Highly expressed in cerebellum, but also found in frontal cortex, hippocampus and basal ganglia.,

Background

This gene encodes a member of a high affinity glutamate transporter family. This gene functions in the termination of excitatory neurotransmission in central nervous system. Mutations are associated with episodic ataxia, Type 6. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Feb 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using EAAT1 Monoclonal Antibody