



EDG-6 Monoclonal Antibody

Catalog No	YP-mAb-13207
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	S1PR4
Immunogen	The antiserum was produced against synthesized peptide derived from human EDG6. AA range:211-260
Specificity	EDG-6 Monoclonal Antibody detects endogenous levels of EDG-6 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EDG-6;S1PR4;Sphingosine 1-phosphate receptor 4;EDG6;S1P receptor 4;S1P4; Endothelial differentiation G-protein coupled receptor 6;Sphingosine 1-phosphate receptor Edg-6;S1P receptor Edg-6
Observed Band	41kD
Calculated Molecular Weight	41kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Specifically expressed in fetal and adult lymphoid and hematopoietic tissue as well as in lung. Considerable level of expression in adult and fetal spleen as well as adult peripheral leukocytes and lung. Lower expression in adult thymus, lymph node, bone marrow, and appendix as well as in fetal liver, thymus, and lung.
Function	function:Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. May be involved in cell migration processes that are specific for lymphocytes.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Specifically expressed in fetal and adult lymphoid and hematopoietic tissue as well as in lung. Considerable level of expression in adult



and fetal spleen as well as adult peripheral leukocytes and lung. Lower expression in adult thymus, lymph node, bone marrow, and appendix as well as in fetal liver, thymus, and lung.,

Background

This gene is a member of the endothelial differentiation, G-protein-coupled (EDG)) receptor gene family. EDG receptors bind lysophospholipids or lysosphingolipids as ligands, and are involved in cell signalling in many different cell types. This EDG receptor gene is intronless and is specifically expressed in the lymphoid tissue. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using EDG-6 Monoclonal Antibody