



IL-4Rα Monoclonal Antibody

Catalog No	YP-mAb-13380
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	IL4R
Immunogen	The antiserum was produced against synthesized peptide derived from human IL-4R/CD124. AA range:463-512
Specificity	IL-4Rα Monoclonal Antibody detects endogenous levels of IL-4Rα protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CD124;IL4RA;IL-4RA;IL4R;IL-4R/CD124;Interleukin-4 receptor subunit alpha; 582J2.1;IL-4 receptor subunit alpha;IL-4R subunit alpha;IL-4R-alpha;CD antigen CD124
Observed Band	90kD
Calculated Molecular Weight	90kD
Cell Pathway	Cell membrane; Single-pass type I membrane protein.; [Isoform 2]: Secreted.
Tissue Specificity	Isoform 1 and isoform 2 are highly expressed in activated T-cells.
Function	domain:Contains 1 copy of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,domain:The box 1 motif is required for JAK interaction and/or activation.,domain:The extracellular domain represents the IL4 binding protein (IL4BP).,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:Receptor for both interleukin 4 and interleukin 13. Couples to the JAK1/2/3-STAT6 pathway. The IL4 response is involved in promoting Th2 differentiation. The IL4/IL13



responses are involved in regulating IgE production and, chemokine and mucus production at sites of allergic inflammation. In certain cel

Background

This gene encodes the alpha chain of the interleukin-4 receptor, a type I transmembrane protein that can bind interleukin 4 and interleukin 13 to regulate IgE production. The encoded protein also can bind interleukin 4 to promote differentiation of Th2 cells. A soluble form of the encoded protein can be produced by proteolysis of the membrane-bound protein, and this soluble form can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells. Allelic variations in this gene have been associated with atopy, a condition that can manifest itself as allergic rhinitis, sinusitis, asthma, or eczema. Polymorphisms in this gene are also associated with resistance to human immunodeficiency virus type-1 infection. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Apr 2012],

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Western Blot analysis of various cells using IL-4Rα Monoclonal Antibody diluted at 1:2000



Western blot analysis of mouse-kidney 293T lysis using IL-4Rα antibody. Antibody was diluted at 1:2000



Western blot analysis of lysates from 293 cells, using IL-4R/CD124 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HUVEC cells using IL-4R/CD124 antibody.