



Nesprin-3 Monoclonal Antibody

Catalog No	YP-mAb-13428
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SYNE3
Immunogen	The antiserum was produced against synthesized peptide derived from human C14orf49. AA range:169-218
Specificity	Nesprin-3 Monoclonal Antibody detects endogenous levels of Nesprin-3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KASH3;NET53;Nesp3;C14orf49;C14orf139;LINC00341;NCRNA00341;SYNE3;Nesprin-3;Nuclear envelope spectrin repeat protein 3
Observed Band	115kD
Calculated Molecular Weight	115kD
Cell Pathway	Nucleus outer membrane ; Single-pass type IV membrane protein . Nucleus envelope . Rough endoplasmic reticulum .
Tissue Specificity	Expressed in aortic endothelial cells (at protein level).
Function	function:Involved in the maintenance of nuclear organization and structural integrity. Probable anchoring protein which theters the nucleus to the cytoskeleton by binding PLEC1/Plectin which can associate with the intermediate filament system.,similarity:Belongs to the nesprin family.,similarity:Contains 1 KASH domain.,similarity:Contains 2 spectrin repeats.,subcellular location:Located at the rough reticulum endoplasmic (RER) in some cells.,subunit:Interacts with actin-binding domain of PLEC1/plectin.,



Background

function:Involved in the maintenance of nuclear organization and structural integrity. Probable anchoring protein which theters the nucleus to the cytoskeleton by binding PLEC1/Plectin which can associate with the intermediate filament system.,similarity:Belongs to the nesprin family.,similarity:Contains 1 KASH domain.,similarity:Contains 2 spectrin repeats.,subcellular location:Located at the rough reticulum endoplasmic (RER) in some cells.,subunit:Interacts with actin-binding domain of PLEC1/plectin.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Nesprin-3 Monoclonal Antibody