



Olfactory receptor 2A4/7 Monoclonal Antibody

Catalog No	YP-mAb-13479
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	OR2A4/OR2A7
Immunogen	The antiserum was produced against synthesized peptide derived from human OR2A4/2A7. AA range:231-280
Specificity	Olfactory receptor 2A4/7 Monoclonal Antibody detects endogenous levels of Olfactory receptor 2A4/7 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Olfactory receptor 2A4/7;OR2A4/OR2A7;OR2A7;Olfactory receptor 2A7; Olfactory receptor OR7-18;OR2A4;OR2A10;Olfactory receptor 2A4;Olfactory receptor 2A10;Olfactory receptor OR6-37
Observed Band	35kD
Calculated Molecular Weight	35kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Liver,
Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
Background	olfactory receptor family 2 subfamily A member 7(OR2A7) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone



receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Olfactory receptor 2A4/7 Monoclonal Antibody