



CD2BP2 Monoclonal Antibody

Catalog No	YP-mAb-13891
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CD2BP2
Immunogen	The antiserum was produced against synthesized peptide derived from human CD2 Tail-binding. AA range:101-150
Specificity	CD2BP2 Monoclonal Antibody detects endogenous levels of CD2BP2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CD2 antigen cytoplasmic tail-binding protein 2;CD2 cytoplasmic domain-binding protein 2;CD2;CD2BP2;KIAA1178;CD2 tail-binding protein 2
Observed Band	50kD
Calculated Molecular Weight	50kD
Cell Pathway	Cytoplasm. Nucleus. Predominantly nuclear.
Tissue Specificity	Aorta,Brain,Epithelium,Liver,Lung,
Function	similarity:Contains 1 GYF domain.,subunit:Binds the cytoplasmic domain of CD2 through the GYF domain.,
Background	This gene encodes a bi-functional protein. In the cytoplasm, the encoded protein binds the cytoplasmic tail of human surface antigen CD2 via its C-terminal GYF domain, and regulate CD2-triggered T lymphocyte activation. In the nucleus, this protein is a component of the U5 small nuclear ribonucleoprotein complex and is involved in RNA splicing. A pseudogene has been identified on chromosome 7. Alternative splicing results in multiple transcript variants but their biological validity has not been determined. [provided by RefSeq, Nov 2008],




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CD2BP2 Monoclonal Antibody