



CD88 Monoclonal Antibody

Catalog No	YP-mAb-13915
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	C5AR1
Immunogen	The antiserum was produced against synthesized peptide derived from human N ternal CD88/C5aR. AA range:301-350
Specificity	CD88 Monoclonal Antibody detects endogenous levels of CD88 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	C5AR1;C5A;C5a anaphylatoxin chemotactic receptor 1;C5a R;C5a-R;CD88;C5a anaphylatoxin chemotactic receptor;C5AR;C5R1;CD antigen CD88
Observed Band	45kD
Calculated Molecular Weight	45kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Cytoplasmic vesicle . Phosphorylated C5aR colocalizes with ARRB1 and ARRB2 in cytoplasmic vesicles. .
Tissue Specificity	Brain,
Function	function:Receptor for the chemotactic and inflammatory peptide anaphylatoxin C5a. This receptor stimulates chemotaxis, granule enzyme release and superoxide anion production.,PTM:Sulfation plays a critical role in the association of the receptor with C5a, but no significant role in the ability of the receptor to transduce a signal and mobilize calcium in response to a small peptide agonist.,similarity:Belongs to the G-protein coupled receptor 1 family.,



Background

function: Receptor for the chemotactic and inflammatory peptide anaphylatoxin C5a. This receptor stimulates chemotaxis, granule enzyme release and superoxide anion production. PTM: Sulfation plays a critical role in the association of the receptor with C5a, but no significant role in the ability of the receptor to transduce a signal and mobilize calcium in response to a small peptide agonist. similarity: Belongs to the G-protein coupled receptor 1 family.


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CD88 Monoclonal Antibody