



Fgl2 Monoclonal Antibody


Catalog No	YP-mAb-13927
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	FGL2
Immunogen	The antiserum was produced against synthesized peptide derived from human Fgl2. AA range:38-87
Specificity	Fgl2 Monoclonal Antibody detects endogenous levels of Fgl2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FGL-2;fibrinogen like 2;Fibrinogen like protein 2;Fibrinogen-like protein 2; Fibroleukin;Fgl2;pT49
Observed Band	55kD
Calculated Molecular Weight	55kD
Cell Pathway	Secreted.
Tissue Specificity	Constitutively expressed in cytotoxic T-cells.
Function	function:May play a role in physiologic lymphocyte functions at mucosal sites., similarity:Contains 1 fibrinogen C-terminal domain.,subunit:Homotetramer; disulfide-linked.,tissue specificity:Constitutively expressed in cytotoxic T-cells.,
Background	fibrinogen like 2(FGL2) Homo sapiens The protein encoded by this gene is a secreted protein that is similar to the beta- and gamma-chains of fibrinogen. The carboxyl-terminus of the encoded protein consists of the fibrinogen-related domains (FRED). The encoded protein forms a tetrameric complex which is stabilized by interchain disulfide bonds. This protein may play a role in physiologic functions at mucosal sites. [provided by RefSeq, Jul 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using Fgl2 Monoclonal Antibody