



CD48 Monoclonal Antibody

Catalog No	YP-mAb-13987
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CD48
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human CD48. AA range:41-90
Specificity	CD48 Monoclonal Antibody detects endogenous levels of CD48 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	BLAST;BLAST1;B-lymphocyte activation marker BLAST-1;CD48 antigen;CD48 molecule;CD48;BCM1;BCM1 surface antigen;Leukocyte antigen MEM-102; TCT.1
Observed Band	40kD
Calculated Molecular Weight	40kD
Cell Pathway	Cell membrane ; Lipid-anchor, GPI-anchor . Secreted .
Tissue Specificity	Widely expressed on all hematopoietic cells.
Function	function:Ligand for CD2. Might facilitate interaction between activated lymphocytes. Probably involved in regulating T-cell activation.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,
Background	This gene encodes a member of the CD2 subfamily of immunoglobulin-like receptors which includes SLAM (signaling lymphocyte activation molecules) proteins. The encoded protein is found on the surface of lymphocytes and other immune cells, dendritic cells and endothelial cells, and participates in activation and differentiation pathways in these cells. The encoded protein does not have a



transmembrane domain, however, but is held at the cell surface by a GPI anchor via a C-terminal domain which maybe cleaved to yield a soluble form of the receptor. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CD48 Monoclonal Antibody