



CD329 Monoclonal Antibody

Catalog No	YP-mAb-14007
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SIGLEC8
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human SIGLEC8. AA range:81-130
Specificity	CD329 Monoclonal Antibody detects endogenous levels of CD329 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SIGLEC9;CD329;Protein FOAP 9;Sialic acid-binding Ig-like lectin 9;Siglec 9; SIGLEC8;Sialic acid-binding Ig-like lectin 8;SAF2;Siglec-8;CDw329; Sialoadhesin family member 2;SAF-2
Observed Band	50kD
Calculated Molecular Weight	50kD
Cell Pathway	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Expressed specifically on red blood cells namely basophil, mast cells and eosinophils.
Function	domain:Contains 1 copy of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,function:Putative adhesion molecule that mediates sialic-acid dependent binding to cells. Preferentially binds to alpha-2,3-linked sialic acid. Also binds to alpha-2,6-linked sialic acid. The sialic acid recognition site may be masked by cis interactions with sialic acids on the same cell surface.,online information:Siglec-8, similarity:Belongs to the immunoglobulin superfamily. SIGLEC (sialic acid



binding Ig-like lectin) family.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Expressed specifically on eosin

Background

Sialic acid-binding immunoglobulin (Ig)-like lectins, or SIGLECs (e.g., CD33 (MIM 159590)), are a family of type 1 transmembrane proteins each having a unique expression pattern, mostly in hemopoietic cells. SIGLEC8 is a member of the CD33-like subgroup of SIGLECs, which are localized to 19q13.3-q13.4 and have 2 conserved cytoplasmic tyrosine-based motifs: an immunoreceptor tyrosine-based inhibitory motif, or ITIM (see MIM 604964), and a motif homologous to one identified in signaling lymphocyte activation molecule (SLAM; MIM 603492) that mediates an association with SLAM-associated protein (SAP; MIM 300490) (summarized by Foussias et al., 2000 [PubMed 11095983]) .[supplied by OMIM, May 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CD329 Monoclonal Antibody