



# FAK Monoclonal Antibody

|                                    |                                                                                                                                                                                                                                                                                                                                                                                             |
|------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Catalog No</b>                  | YP-mAb-14745                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Isotype</b>                     | IgG                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Reactivity</b>                  | Human;Mouse;Rat                                                                                                                                                                                                                                                                                                                                                                             |
| <b>Applications</b>                | WB                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Gene Name</b>                   | PTK2                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Immunogen</b>                   | The antiserum was produced against synthesized peptide derived from human FAK. AA range:809-858                                                                                                                                                                                                                                                                                             |
| <b>Specificity</b>                 | FAK Monoclonal Antibody detects endogenous levels of FAK protein.                                                                                                                                                                                                                                                                                                                           |
| <b>Formulation</b>                 | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                                                                                                                                                                                                     |
| <b>Source</b>                      | Monoclonal, Mouse,IgG                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Purification</b>                | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.                                                                                                                                                                                                                                                                        |
| <b>Dilution</b>                    | WB 1:500-1:2000                                                                                                                                                                                                                                                                                                                                                                             |
| <b>Concentration</b>               | 1 mg/ml                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Purity</b>                      | ≥90%                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Storage Stability</b>           | -20°C/1 year                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Synonyms</b>                    | PTK2;EC:2.7.10.2;FADK 1;FAK1;Focal adhesion kinase 1;FAK;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71; PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK                                                                                                                                                                                       |
| <b>Observed Band</b>               | 120kD                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Calculated Molecular Weight</b> | 120kD                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Cell Pathway</b>                | Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Nucleus. Cytoplasm, cytoskeleton, cilium basal body . Constituent of focal adhesions. Detected at microtubules.                                                          |
| <b>Tissue Specificity</b>          | Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787).                                                                                                                                                                                               |
| <b>Function</b>                    | catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,domain:The carboxy-terminal region is the site of focal adhesion targeting (FAT) sequence which mediates the localization of FAK1 to focal adhesions.,domain:The first Pro-rich domain interacts with the SH3 domain of CRK-associated substrate (BCAR1) and CASL.,function:Non-receptor protein- |



tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity., PTM:Phosphorylated on 6 tyrosine residues upon activation

**Background**

protein tyrosine kinase 2(PTK2) Homo sapiens This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. Several transcript variants encoding different isoforms have been found for this gene, but the full-length nature of only four of them have been determined. [provided by RefSeq, Oct 2015],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images