



GRK 1 Monoclonal Antibody

Catalog No	YP-mAb-14762
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	GRK1
Immunogen	The antiserum was produced against synthesized peptide derived from human GRK1. AA range:6-55
Specificity	GRK 1 Monoclonal Antibody detects endogenous levels of GRK 1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EC:2.7.11.14;G protein-coupled receptor kinase 1;GPRK1;Rhodopsin kinase; Rhodopsin kinase GRK1;GRK1;RHOK;RK
Observed Band	63kD
Calculated Molecular Weight	63kD
Cell Pathway	Membrane ; Lipid-anchor . Cell projection, cilium, photoreceptor outer segment . Subcellular location is not affected by light or dark conditions. .
Tissue Specificity	Retinal-specific. Expressed in rods and cones cells.
Function	catalytic activity:ATP + [rhodopsin] = ADP + [rhodopsin] phosphate., disease:Defects in GRK1 are a cause of congenital stationary night blindness Oguchi type (CSNBO) [MIM:258100]; also known as Oguchi disease. Congenital stationary night blindness is a non-progressive retinal disorder characterized by impaired night vision. CSNBO is an autosomal recessive form associated with fundus discoloration and abnormally slow dark adaptation., function:Phosphorylates rhodopsin thereby initiating its deactivation.,online information:Retina International's Scientific Newsletter, PTM:Autophosphorylated.,PTM:Farnesylation is required for full activity., similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase



family. GPRK subfamily.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 RGS domain.,tissue specificity:R

Background

This gene encodes a member of the guanine nucleotide-binding protein (G protein)-coupled receptor kinase subfamily of the Ser/Thr protein kinase family. The protein phosphorylates rhodopsin and initiates its deactivation. Defects in GRK1 are known to cause Oguchi disease 2 (also known as stationary night blindness Oguchi type-2). [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using GRK 1 Monoclonal Antibody