



HGK Monoclonal Antibody

Catalog No	YP-mAb-14773
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	MAP4K4
Immunogen	The antiserum was produced against synthesized peptide derived from human MEKKK 4. AA range:406-455
Specificity	HGK Monoclonal Antibody detects endogenous levels of HGK protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EC:2.7.11.1;FLH21957;HGK;HPK/GCK like kinase HGK;HPK/GCK-like kinase HGK;MAP4K4;Mitogen-activated protein kinase kinase kinase kinase 4; KIAA0687;NIK;MAPK/ERK kinase kinase kinase 4;MEK kinase kinase 4;MEKKK 4;Nck-interacting kinase
Observed Band	142kD
Calculated Molecular Weight	142kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Widely expressed. Isoform 5 is abundant in the brain. Isoform 4 is predominant in the liver, skeletal muscle and placenta.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein., cofactor:Magnesium.,function:Serine/threonine kinase that may play a role in the response to environmental stress and cytokines such as TNF-alpha. Appears to act upstream of the JUN N-terminal pathway.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily.,similarity:Contains 1 CNH domain., similarity:Contains 1 protein kinase domain.,subunit:Interacts with the SH3



domain of the adapter proteins Nck (By similarity). Binds, via its CNH regulatory domain, to the N-terminal region of SPG3A.,tissue specificity:Appears to be ubiquitous, expressed in all tissue types examined. Isoform 5 appears to be more abundant in the brain, isoform 4 is predominant in the liver, skelet

Background

mitogen-activated protein kinase kinase kinase 4(MAP4K4) Homo sapiens The protein encoded by this gene is a member of the serine/threonine protein kinase family. This kinase has been shown to specifically activate MAPK8/JNK. The activation of MAPK8 by this kinase is found to be inhibited by the dominant-negative mutants of MAP3K7/TAK1, MAP2K4/MKK4, and MAP2K7/MKK7, which suggests that this kinase may function through the MAP3K7-MAP2K4-MAP2K7 kinase cascade, and mediate the TNF-alpha signaling pathway. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using HGK Monoclonal Antibody