



MARK1/2/3/4 Monoclonal Antibody


Catalog No	YP-mAb-14824
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	MARK1/2/3/4
Immunogen	The antiserum was produced against synthesized peptide derived from human MARK1/2/3/4. AA range:181-230
Specificity	MARK1/2/3/4 Monoclonal Antibody detects endogenous levels of MARK1/2/3/4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KIAA1477;MAP/microtubule affinity regulating kinase 1;MAP/microtubule affinity-regulating kinase 1;MARK 1;MARK;Mark1;MARK1_HUMAN;MGC126512;MGC126513;Serine/threonine protein kinase MARK1;Serine/threonine-protein kinase MARK1;ELKL motif kinase 1;ELKL motif kinase;EMK-1;EMK1;MAP/microtubule affinity regulating kinase 2;MAP/microtubule affinity-regulating kinase 2;Mark2;MARK2_HUMAN;MGC99619;PAR 1;Par 1b;PAR1 homolog;Par1b;Ser/Thr protein kinase PAR 1B;Serine/threonine protein kinase EMK;Serine/threonine protein kinase MARK2;Serine/threonine-protein kinase MARK2;C-TAK1;Cdc25C associated protein kinase 1;Cdc25C-associated protein kinase 1;cTAK1;ELKL motif kinase 2;EMK-2;Emk2;ETK 1;KIAA423;KP78;MAP microtubule affinity regulating kinase 3;MAP/microtubule affinity-regulating kinase 3;Mark3;MARK3_HUMAN;Par 1a;PAR1A;Protein kinase STK1;Ser/Thr protein kinase PAR-1;Serine threonine protein kinase p78;Serine/threonine-protein kinase p78;SerThr protein kinase PAR 1;FLJ997;KIAA186;MAP / microtubule affinity regulating kinase 4;MAP / microtubule affinity regulating kinase 4L;MAP/microtubule affinity regulating kinase 4;MAP/microtubule affinity regulating kinase like 1;MAP/microtubule affinity-regulating kinase 4;MAP/microtubule affinity-regulating kinase-like 1;MARK 4;MARK 4L;MARK L1;MARK4;MARK4 serine / threonine protein kinase;MARK4 serine/threonine protein kinase;MARK4_HUMAN;MARK4S;MARKL 1;MARKL1;Nbla65;Putative protein product of Nbla65;MARK1/2/3/4;Serine/threonine-protein kinase MARK1/2/3/4;PAR1 homolog c;Par-1c;Par1c;MAP/microtubule affi



Observed Band	89kD
Calculated Molecular Weight	89kD
Cell Pathway	Cell membrane ; Peripheral membrane protein . Cytoplasm, cytoskeleton . Cytoplasm . Cell projection, dendrite . Appears to localize to an intracellular network. .
Tissue Specificity	Highly expressed in heart, skeletal muscle, brain, fetal brain and fetal kidney.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein., cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Thr-215 by STK11 in complex with STE20-related adapter-alpha (STRAD alpha) pseudo kinase and CAB39.,function:May play a role in cytoskeletal stability., similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. MARK subfamily.,similarity:Contains 1 KA1 (kinase-associated) domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 UBA domain.,subcellular location:Appears to localize to an intracellular network., tissue specificity:Highly expressed in heart, skeletal muscle, brain, fetal brain and fetal kidney.,
Background	catalytic activity:ATP + a protein = ADP + a phosphoprotein., cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Thr-215 by STK11 in complex with STE20-related adapter-alpha (STRAD alpha) pseudo kinase and CAB39.,function:May play a role in cytoskeletal stability., similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. MARK subfamily.,similarity:Contains 1 KA1 (kinase-associated) domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 UBA domain.,subcellular location:Appears to localize to an intracellular network., tissue specificity:Highly expressed in heart, skeletal muscle, brain, fetal brain and fetal kidney.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using
MARK1/2/3/4 Monoclonal Antibody