



MEK-4 Monoclonal Antibody

Catalog No	YP-mAb-14839
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	MAP2K4
Immunogen	The antiserum was produced against synthesized peptide derived from human SEK1/MKK4. AA range:46-95
Specificity	MEK-4 Monoclonal Antibody detects endogenous levels of MEK-4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MEK4;c-Jun N-terminal kinase kinase 1;Dual specificity mitogen-activated protein kinase kinase 4;EC:2.7.12.2;JNK activating kinase 1;MEK-4;MAP2K4;JNKK1;MKK4;PRKMK4;SEK1;SERK1;SKK1;MAP kinase kinase 4;MAPKK 4;JNK-activating kinase 1;MAPK/ERK kinase 4;MEK 4;SAPK/ERK kinase 1;Stress-activated pro
Observed Band	44kD
Calculated Molecular Weight	44kD
Cell Pathway	Cytoplasm . Nucleus .
Tissue Specificity	Abundant expression is seen in the skeletal muscle. It is also widely expressed in other tissues.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Dual specificity kinase that activates the JUN kinases MAPK8 (JNK1) and MAPK9 (JNK2) as well as MAPK14 (p38) but not MAPK1 (ERK2) or MAPK3 (ERK1)., PTM:Activated by phosphorylation on Ser/Thr by MAP kinase kinase kinases., similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Interacts



with SPAG9.,tissue specificity:Abundant expression is seen in the skeletal muscle. It is also widely expressed in other tissues.,

Background

This gene encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this family act as an integration point for multiple biochemical signals and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKKs, MAPKKs, and MAPKs. This protein is phosphorylated at serine and threonine residues by MAPKKKs and subsequently phosphorylates downstream MAPK targets at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver organogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using MEK-4 Monoclonal Antibody