



Meis1 Monoclonal Antibody

Catalog No	YP-mAb-15777
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	MEIS1
Immunogen	The antiserum was produced against synthesized peptide derived from human Meis1. AA range:229-278
Specificity	Meis1 Monoclonal Antibody detects endogenous levels of Meis1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Homeobox protein Meis1;Meis1
Observed Band	37kD
Calculated Molecular Weight	37kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed at low level in normal immunohepatopoietic tissues, including the fetal liver. Expressed in a subset of myeloid leukemia cell lines, with the highest expression seen in those with a megakaryocytic-erythroid phenotype. Also expressed at high levels in the cerebellum.
Function	function:Acts as a transcriptional regulator of PAX6. Acts as a transcriptional activator of PF4 in complex with PBX1 or PBX2. Required for hematopoiesis, megakaryocyte lineage development and vascular patterning. May function as a cofactor for HOXA7 and HOXA9 in the induction of myeloid leukemias., similarity:Belongs to the TALE/MEIS homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,subunit:Interacts with the N-terminal region of PBX1 to form a heterodimer which binds DNA including a cAMP-responsive sequence in CYP17. Also forms heterodimers with PBX2. Forms heterotrimers with PBX1 or PBX2 and a number of HOX proteins including HOXA9, HOXD4



and HOXD9 where it acts as a non-DNA-binding partner. Also forms heterotrimers with PBX1 and HOX proteins including HOXD9 and HOXD10 where PBX1 is the non-DNA-binding partner.,tissue specificity:Expressed at low level in normal immun

Background

Homeobox genes, of which the most well-characterized category is represented by the HOX genes, play a crucial role in normal development. In addition, several homeoproteins are involved in neoplasia. This gene encodes a homeobox protein belonging to the TALE (‘three amino acid loop extension’) family of homeodomain-containing proteins. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Meis1 Monoclonal Antibody