



Cleaved-Ephrin-A2 (N188) Monoclonal Antibody

Catalog No	YP-mAb-15812
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	EFNA2
Immunogen	The antiserum was produced against synthesized peptide derived from human EFNA2. AA range:139-188
Specificity	Cleaved-Ephrin-A2 (N188) Monoclonal Antibody detects endogenous levels of fragment of activated Ephrin-A2 protein resulting from cleavage adjacent to N188.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EFNA 2;efna2;EFNA2_HUMAN;ELF1;Eph related receptor tyrosine kinase ligand 6;EPH-related receptor tyrosine kinase ligand 6;Ephrin-A2;EphrinA2;EPLG6;HEK7 L;HEK7 ligand;HEK7-L;HEK7L;LERK 6;LERK-6;LERK6;Ligand of eph related kinase 6
Observed Band	21kD
Calculated Molecular Weight	21kD
Cell Pathway	Cell membrane ; Lipid-anchor, GPI-anchor .
Tissue Specificity	Brain,
Function	similarity:Belongs to the ephrin family.,subunit:Binds to the receptor tyrosine kinases EPHA3, EPHA4 and EPHA5.,
Background	This gene encodes a member of the ephrin family. The protein is composed of a signal sequence, a receptor-binding region, a spacer region, and a hydrophobic region. The EPH and EPH-related receptors comprise the largest subfamily of receptor protein-tyrosine kinases and have been implicated in mediating



developmental events, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. Posttranslational modifications determine whether this protein localizes to the nucleus or the cytoplasm. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Cleaved-Ephrin-A2 (N188) Monoclonal Antibody