



Cleaved-ITI-H2 (D702) Monoclonal Antibody

Catalog No	YP-mAb-15816
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ITIH2
Immunogen	The antiserum was produced against synthesized peptide derived from human ITIH2. AA range:653-702
Specificity	Cleaved-ITI-H2 (D702) Monoclonal Antibody detects endogenous levels of fragment of activated ITI-H2 protein resulting from cleavage adjacent to D702.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AI74722;H2P;IGHEP2;Inter alpha (globulin) inhibitor H2;Inter alpha globulin inhibitor H2 polypeptide;Inter alpha trypsin inhibitor complex component II;Inter alpha trypsin inhibitor heavy chain H2;Inter-alpha-inhibitor heavy chain 2;Inter-alpha-trypsin inhibitor complex component II;Inter-alpha-trypsin inhibitor heavy chain H2;Intin2;ITI HC2;ITI heavy chain H2;ITI-HC2;Itih 2;Itih2;ITIH2_HUMAN; OTTHUMP1958;OTTHUMP1959;OTTMUSP1219;RP23 119N4.3;Serum derived hyaluronan associated protein;Serum-derived hyaluronan-associated protein; SHAP;ITI-H2
Observed Band	72kD
Calculated Molecular Weight	72kD
Cell Pathway	Secreted.
Tissue Specificity	Plasma.
Function	function:May act as a carrier of hyaluronan in serum or as a binding protein between hyaluronan and other matrix protein, including those on cell surfaces in tissues to regulate the localization, synthesis and degradation of hyaluronan which are essential to cells undergoing biological processes.,mass



spectrometry: PubMed:9677337,PTM:Heavy chains are linked to bikunin via chondroitin 4-sulfate esterified to the alpha-carboxyl of the C-terminal aspartate after propeptide cleavage.,similarity:Belongs to the ITIH family.,similarity:Contains 1 VWFA domain.,subunit:I-alpha-I plasma protease inhibitors are assembled from one or two heavy chains (H1, H2 or H3) and one light chain, bikunin. Inter-alpha-inhibitor (I-alpha-I) is composed of H1, H2 and bikunin, inter-alpha-like inhibitor (I-alpha-LI) of H2 and bikunin, and pre-alpha-inhibitor (P-alpha-I) of H3 and bikunin.,

Background

The inter-alpha-trypsin inhibitors (ITI) are a family of structurally related plasma serine protease inhibitors involved in extracellular matrix stabilization and in prevention of tumor metastasis. The ITI family contains multiple proteins made up of a light chain (see MIM 176870) and a variable number of heavy chains (Salier et al., 1987 [PubMed 2446322]; Himmelfarb et al., 2004 [PubMed 14744536]).[supplied by OMIM, Nov 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Cleaved-ITI-H2 (D702) Monoclonal Antibody