



Eotaxin-3 Monoclonal Antibody

Catalog No	YP-mAb-15989
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CCL26
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human CCL26. AA range:31-80
Specificity	Eotaxin-3 Monoclonal Antibody detects endogenous levels of Eotaxin-3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	C-C motif chemokine 26;CC chemokine IMAC;CCL 26;CCL26;CCL26_HUMAN; Chemokine (C C motif) ligand 26;chemokine N1;Eotaxin-3;IMAC;Macrophage inflammatory protein 4-alpha;MIP 4a;MIP 4alpha;MIP-4-alpha;MIP4a;MIP4alpha; SCYA 26;SCYA26;Small inducible cytokine A26;Small inducible cytokine subfamily A (Cys Cys) member 26;Small inducible cytokine subfamily A member 26;Small-inducible cytokine A26;Thymic stroma chemokine 1;Thymic stroma chemokine-1;TSC 1;TSC-1;TSC1
Observed Band	12kD
Calculated Molecular Weight	12kD
Cell Pathway	Secreted .
Tissue Specificity	Ubiquitously expressed at low levels in various tissues including heart and ovary.
Function	function:Chemotactic for eosinophils and basophils. Binds to CCR3.,online information:CCL26 entry,similarity:Belongs to the intercrine beta (chemokine CC) family.,subunit:Monomer.,tissue specificity:Ubiquitously expressed at low levels in various tissues including heart and ovary.,



Background

This gene is one of two Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 7. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for normal peripheral blood eosinophils and basophils. The product of this gene is one of three related chemokines that specifically activate chemokine receptor CCR3. This chemokine may contribute to the eosinophil accumulation in atopic diseases. [provided by RefSeq, Jul 2008]


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Eotaxin-3 Monoclonal Antibody