



Neuregulin-2 Monoclonal Antibody

Catalog No	YP-mAb-16009
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	NRG2
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human NRG2. AA range:361-410
Specificity	Neuregulin-2 Monoclonal Antibody detects endogenous levels of Neuregulin-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	neuregulin 2;neuregulin2;Neuregulin-2;NTAK;Pro NRG2;NRG2;Pro-neuregulin-2 membrane-bound isoform;Pro-neuregulin-2;membrane-bound isoform;Pro-NRG2; membrane-bound isoform
Observed Band	90kD
Calculated Molecular Weight	90kD
Cell Pathway	[Pro-neuregulin-2, membrane-bound isoform]: Cell membrane ; Single-pass type I membrane protein . Does not seem to be active. .; [Neuregulin-2]: Secreted .
Tissue Specificity	Restricted to the cerebellum in the adult.
Function	domain:ERBB receptor binding is elicited entirely by the EGF-like domain., domain:The cytoplasmic domain may be involved in the regulation of trafficking and proteolytic processing. Regulation of the proteolytic processing involves initial intracellular domain dimerization.,function:Direct ligand for ERBB3 and ERBB4 tyrosine kinase receptors. Concomitantly recruits ERBB1 and ERBB2 coreceptors, resulting in ligand-stimulated tyrosine phosphorylation and activation of the ERBB receptors. May also promote the heterodimerization with the EGF receptor.,PTM:Extensive glycosylation precedes the proteolytic cleavage.,PTM:Proteolytic cleavage close to the plasma membrane on the



external face leads to the release of the soluble growth factor form., similarity:Belongs to the neuregulin family.,similarity:Contains 1 EGF-like domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,s

Background

This gene encodes a novel member of the neuregulin family of growth and differentiation factors. Through interaction with the ERBB family of receptors, this protein induces the growth and differentiation of epithelial, neuronal, glial, and other types of cells. The gene consists of 12 exons and the genomic structure is similar to that of neuregulin 1, another member of the neuregulin family of ligands. The products of these genes mediate distinct biological processes by acting at different sites in tissues and eliciting different biological responses in cells. This gene is located close to the region for demyelinating Charcot-Marie-Tooth disease locus, but is not responsible for this disease. Alternative transcript variants encoding distinct isoforms have been described. [provided by RefSeq, May 2010],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Neuregulin-2 Monoclonal Antibody