



TPO Monoclonal Antibody

Catalog No	YP-mAb-16033
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	THPO
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human THPO. AA range:41-90
Specificity	TPO Monoclonal Antibody detects endogenous levels of TPO protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MSA;PERT_HUMAN;TDH2A;Thyroid microsomal antigen;Thyroid peroxidase;Thyropoxidase;TPO;TPX;THPO;Thrombopoietin;MGDF;C-mpl ligand;ML;Megakaryocyte colony-stimulating factor;Megakaryocyte growth and development factor;Myeloproliferative leukemia virus oncogene ligand
Observed Band	38kD
Calculated Molecular Weight	38kD
Cell Pathway	Secreted.
Tissue Specificity	Brain,Fetal liver,Liver,Placenta,
Function	disease:Defects in THPO are a cause of essential thrombocythemia (ET) [MIM:187950]. ET is inherited as an autosomal dominant trait which is characterized by elevated platelet levels due to sustained proliferation of megakaryocytes, and frequently lead to thrombotic and haemorrhagic complications.,domain:Two-domain structure with an erythropoietin-like N-terminal and a Ser/Pro/Thr-rich C-terminal.,function:Lineage-specific cytokine affecting the proliferation and maturation of megakaryocytes from their committed progenitor cells. It acts at a late stage of megakaryocyte development. It may be the major physiological regulator of circulating platelets.,



similarity:Belongs to the EPO/TPO family.,

Background

Megakaryocytopoiesis is the cellular development process that leads to platelet production. The main functional protein encoded by this gene is a humoral growth factor that is necessary for megakaryocyte proliferation and maturation, as well as for thrombopoiesis. This protein is the ligand for MLP/C_MPL, the product of myeloproliferative leukemia virus oncogene. Mutations in this gene are the cause of thrombocythemia 1. Alternative promoter usage and differential splicing result in multiple transcript variants differing in the 5' UTR and/or coding region. Multiple AUG codons upstream of the main open reading frame (ORF) have been identified, and these upstream AUGs inhibit translation of the main ORF at different extent. [provided by RefSeq, Feb 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TPO Monoclonal Antibody