



FGF-17 Monoclonal Antibody

Catalog No	YP-mAb-16054
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	FGF17
Immunogen	The antiserum was produced against synthesized peptide derived from the C-terminal region of human FGF17. AA range:267-216
Specificity	FGF-17 Monoclonal Antibody detects endogenous levels of FGF-17 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FGF 17;FGF-17;Fibroblast growth factor 17;UNQ161/PRO187;FGF17
Observed Band	25kD
Calculated Molecular Weight	25kD
Cell Pathway	Secreted.
Tissue Specificity	Preferentially expressed in the embryonic brain.
Function	developmental stage:Detected in embryos at E14.5, but not at E10.5 and E19.5. Preferentially expressed in the neuroepithelia of the isthmus and septum of the embryonic brain at E14.5.,function:May be a signaling molecule in the induction and patterning of the embryonic brain.,similarity:Belongs to the heparin-binding growth factors family.,tissue specificity:Preferentially expressed in the embryonic brain.,
Background	This gene encodes a member of the fibroblast growth factor (FGF) family. Member of the FGF family possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes including embryonic development cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein is expressed during embryogenesis and in the adult



cerebellum and cortex and may be essential for vascular growth and normal brain development. Mutations in this gene are the cause of hypogonadotropic hypogonadism 20 with or without anosmia. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Jan 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using FGF-17 Monoclonal Antibody