




IL-1F5 Monoclonal Antibody

Catalog No	YP-mAb-16073
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	IL36RN
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human IL36RN. AA range:1-50
Specificity	IL-1F5 Monoclonal Antibody detects endogenous levels of IL-1F5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AI413231;Family of interleukin 1 delta;FIL1;FIL1 delta;FIL1(DELTA);FIL1D;Fil1delta;I36RA_HUMAN;IL 1 delta;IL 1 related protein 3;IL 1F5 (IL 1HY1;IL 1RP3;IL 1L1;IL 1 delta);il 1h3;IL 1HY1;IL 1ra homolog;IL1F5 (Canonical product IL 1F5a);IL-1 delta;IL-1-related protein 3;IL-1F5;IL-1HY1;IL-1L1;IL-1ra homolog 1;IL-1RP3;IL1F5;IL1HY1;IL1L1;IL1RP3;IL36RA;IL36RN;Interleukin 1 delta; Interleukin 1 family member 5 (delta);Interleukin 1 family member 5;Interleukin 1 HY1;Interleukin 1 like protein 1;Interleukin 1 receptor antagonist homolog 1; Interleukin 36 receptor antagonist;Interleukin-1 delta;Interleukin-1 family member 5;Interleukin-1 HY1;Interleukin-1 receptor antagonist homolog 1; Interleukin-1-like protein 1;Interleukin-36 receptor antagonist protein;MGC2984; PSORP;RP23-176J12.6
Observed Band	17kD
Calculated Molecular Weight	17kD
Cell Pathway	Cytoplasm . Secreted . The secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10; it results in protein translocation from the cytoplasm into the ERGIC (endoplasmic reticulum-Golgi intermediate compartment) followed by vesicle entry and secretion. .



Tissue Specificity	Predominantly expressed in skin keratinocytes but not in fibroblasts, endothelial cells or melanocytes. Detected also in the spleen, brain leukocyte and macrophage cell types. Increased in lesional psoriasis skin.
Function	function:Is a highly and a specific antagonist of the IL-1 receptor-related protein 2-mediated response to interleukin 1 family member 9 (IL1F9). Could constitute part of an independent signaling system analogous to interleukin-1 alpha (IL-1A), beta (IL-1B) receptor agonist and interleukin-1 receptor type I (IL-1R1), that is present in epithelial barriers and takes part in local inflammatory response., induction:By phorbol ester (PMA) and lipopolysaccharide (LPS) treatment in macrophage cell line.,online information:Interleukin-1 entry,similarity:Belongs to the IL-1 family.,tissue specificity:Predominantly expressed in keratinocytes but not in fibroblasts, endothelial cells or melanocytes. Detected also in the spleen, brain leukocyte and macrophage cell types. Increased in lesional psoriasis skin.,
Background	The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine was shown to specifically inhibit the activation of NF-kapMAB induced by interleukin 1 family, member 6 (IL1F6). This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. Two alternatively spliced transcript variants encoding the same protein have been reported. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IL-1F5 Monoclonal Antibody