



TASK-5 Monoclonal Antibody

Catalog No	YP-mAb-16501
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	KCNK15
Immunogen	The antiserum was produced against synthesized peptide derived from human KCNK15. AA range:273-322
Specificity	TASK-5 Monoclonal Antibody detects endogenous levels of TASK-5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Acid sensitive potassium channel protein TASK 5;Acid-sensitive potassium channel protein TASK-5;dJ781B1.1;K2p15.1;KCNK11;KCNK14;Kcnk15;KCNKF_HUMAN;KIAA237;KT3.3;OTTHUMP3172;Potassium channel subfamily K member 14;Potassium channel subfamily K member 15;Potassium family subfamily K member 15;TASK 5;TASK5;TWIK related acid sensitive K(+) channel 5;TWIK-related acid-sensitive K(+) channel 5;Two pore K(+) channel KT3.3;Two pore potassium channel KT3.3;TASK-5
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Membrane; Multi-pass membrane protein.
Tissue Specificity	Detected in pancreas, heart, placenta, lung, liver, kidney, ovary, testis, skeletal muscle and adrenal gland, and at lower levels in prostate, spleen and thyroid gland.
Function	function:Probable potassium channel subunit. No channel activity observed in heterologous systems. May need to associate with another protein to form a functional channel.,polymorphism:Three variant polypeptides are known: TASK-5A, TASK-5B and TASK-5C. The sequence shown is that of TASK-5C.,



similarity:Belongs to the two pore domain potassium channel (TC 1.A.1.8) family.,subunit:Heterodimer .,tissue specificity:Detected in pancreas, heart, placenta, lung, liver, kidney, ovary, testis, skeletal muscle and adrenal gland, and at lower levels in prostate, spleen and thyroid gland.,

Background

potassium two pore domain channel subfamily K member 15(KCNK15) Homo sapiens This gene encodes one of the members of the superfamily of potassium channel proteins containing two pore-forming P domains. The product of this gene has not been shown to be a functional channel, however, it may require other non-pore-forming proteins for activity. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TASK-5 Monoclonal Antibody