



BOCT Monoclonal Antibody


Catalog No	YP-mAb-16519
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SLC22A17
Immunogen	Synthesized peptide derived from the Internal region of human BOCT.
Specificity	BOCT Monoclonal Antibody detects endogenous levels of BOCT protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	24p3 receptor;24p3R;BOCT;BOIT;Brain type organic cation transporter;Brain-type organic cation transporter;hBOIT;Lipocalin-2 receptor;Neutrophil gelatinase associated lipocalin receptor;Neutrophil gelatinase-associated lipocalin receptor;NGAL receptor;NgalR;NGALR2;NGALR3;OTTHUMP8269;OTTHUMP82691; Potent brain type organic ion transporter;S22AH_HUMAN;Slc22a17;Solute carrier family 22 member 17;Solute carrier family 22 organic cation transporter member 17
Observed Band	57kD
Calculated Molecular Weight	57kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Vacuole membrane ; Multi-pass membrane protein . Upon LCN2-binding, it is internalized.
Tissue Specificity	Expressed in brain.
Function	similarity:Belongs to the major facilitator superfamily. Organic cation transporter family.,tissue specificity:Expressed in brain.,
Background	similarity:Belongs to the major facilitator superfamily. Organic cation transporter family.,tissue specificity:Expressed in brain.,

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using BOCT Monoclonal Antibody