



Cdc27 Monoclonal Antibody

Catalog No	YP-mAb-16682
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CDC27
Immunogen	The antiserum was produced against synthesized peptide derived from human H-NUC. AA range:361-410
Specificity	Cdc27 Monoclonal Antibody detects endogenous levels of Cdc27 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CDC27;CDC27 homolog;APC3;APC 3;Anaphase-promoting complex subunit 3; Cell division cycle protein 27 homolog;ANAPC3;D0S1430E;D17S978E; CDC27Hs;H-NUC
Observed Band	92kD
Calculated Molecular Weight	92kD
Cell Pathway	Nucleus . Cytoplasm, cytoskeleton, spindle .
Tissue Specificity	Epithelium,Uterus,
Function	PTM:Phosphorylated. Phosphorylation on Ser-426 and Thr-446 occurs specifically during mitosis.,similarity:Belongs to the APC3/CDC27 family., similarity:Contains 9 TPR repeats.,subunit:Interacts with RB.,
Background	The protein encoded by this gene shares strong similarity with Saccharomyces cerevisiae protein Cdc27, and the gene product of Schizosaccharomyces pombe nuc 2. This protein is a component of the anaphase-promoting complex (APC), which is composed of eight protein subunits and is highly conserved in eukaryotic cells. This complex catalyzes the formation of cyclin B-ubiquitin conjugate, which is responsible for the ubiquitin-mediated proteolysis of B-type



cyclins. The protein encoded by this gene and three other members of the APC complex contain tetratricopeptide (TPR) repeats, which are important for protein-protein interactions. This protein was shown to interact with mitotic checkpoint proteins including Mad2, p55CDC and BUBR1, and it may thus be involved in controlling the timing of mitosis. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been i


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Cdc27 Monoclonal Antibody