Cks2 Mouse mAb

Catalog No	YP-mAb-16713
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	
Immunogen	Recombinant protein (or fragment).This information is considered to be commercially sensitive.
Specificity	
Formulation	
Source	Monoclonal, Mouse,IgG
Purification	Affinity purification
Dilution	WB: 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CDC28;CDC28 protein kinase 2;CDC28 protein kinase regulatory subunit 2;CKS 2;CKS-2;CKS1(S. cerevisiae Cdc28/Cdc2 kinase subunit) homolog 2;Cks2; CKS2_HUMAN;CKSHS2;Cyclin dependent kinases regulatory subunit 2;Cyclin-dependent kinases regulatory subunit 2
Observed Band	14kDa
Calculated Molecular Weight	10kDa
Cell Pathway	
Tissue Specificity	
Function	
Background	CKS2 protein binds to the catalytic subunit of the cyclin dependent kinases and is essential for their biological function. The CKS2 mRNA is found to be expressed in different patterns through the cell cycle in HeLa cells, which reflects specialized role for the encoded protein.
matters needing attention	Avoid repeated freezing and thawing!

Usage suggestions This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from rat liver, using CKS2 Mouse mAb (A3791) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit.
Exposure time: 30s.