



Claudin-4 Monoclonal Antibody

Catalog No	YP-mAb-16959
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CLDN4
Immunogen	The antiserum was produced against synthesized peptide derived from human Claudin 4. AA range:160-209
Specificity	Claudin-4 Monoclonal Antibody detects endogenous levels of Claudin-4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Claudin 4;CLDN4;CPE R;CPE receptor;CPER;Claudin-4;CPETR1;WBSCR8; Clostridium perfringens enterotoxin receptor;CPE-R;CPE-receptor;Williams-Beuren syndrome chromosomal region 8 protein
Observed Band	22kD
Calculated Molecular Weight	22kD
Cell Pathway	Cell junction, tight junction . Cell membrane ; Multi-pass membrane protein . CLDN4 is required for tight junction localization in the kidney. .
Tissue Specificity	Colon,Fetal brain,Trachea,
Function	disease:Haploinsufficiency of CLDN4 may be the cause of certain cardiovascular and musculo-skeletal abnormalities observed in Williams-Beuren syndrome (WBS), a rare developmental disorder. It is a contiguous gene deletion syndrome involving genes from chromosome band 7q11.23., function:Plays a major role in tight junction-specific obliteration of the intercellular space.,similarity:Belongs to the claudin family.,subunit:Directly interacts with TJP1/ZO-1, TJP2/ZO-2 and TJP3/ZO-3.,



Background

The protein encoded by this intronless gene belongs to the claudin family. Claudins are integral membrane proteins that are components of the epithelial cell tight junctions, which regulate movement of solutes and ions through the paracellular space. This protein is a high-affinity receptor for Clostridium perfringens enterotoxin (CPE) and may play a role in internal organ development and function during pre- and postnatal life. This gene is deleted in Williams-Beuren syndrome, a neurodevelopmental disorder affecting multiple systems. [provided by RefSeq, Sep 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Claudin-4 Monoclonal Antibody