



COL11A1 Monoclonal Antibody

Catalog No	YP-mAb-16967
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	COL11A1
Immunogen	The antiserum was produced against synthesized peptide derived from human Collagen XI alpha1. AA range:581-630
Specificity	COL11A1 Monoclonal Antibody detects endogenous levels of COL11A1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	COL11;COL11A1;CO11A1;COLL6;Collagen alpha 1(XI) chain;Collagen XI α1; Collagen alpha-1(XI) chain
Observed Band	181kD
Calculated Molecular Weight	181kD
Cell Pathway	Secreted, extracellular space, extracellular matrix .
Tissue Specificity	Cartilage, placenta and some tumor or virally transformed cell lines. Isoforms using exon IIA or IIB are found in the cartilage while isoforms using only exon IIB are found in the tendon.
Function	alternative products:Additional isoforms seem to exist. There is alternative usage of exon IIA or exon IIB. Transcripts containing exon IIA or IIB are present in cartilage, but exon IIB is preferentially utilized in transcripts from tendon, disease:Defects in COL11A1 are the cause of Marshall syndrome [MIM:154780] . It is an autosomal dominant disorder with ocular, orofacial, auditory and skeletal manifestations. It shares several features with Stickler syndrome, such as midfacial hypoplasia, high myopia, and sensorineural-hearing deficit., disease:Defects in COL11A1 are the cause of Stickler syndrome type 2 (STL2) [MIM:604841]; also known as Stickler syndrome vitreous type 2. STL2 is an



autosomal dominant form of Stickler syndrome, an inherited disorder that associates ocular signs with more or less complete forms of Pierre Robin sequence, bone disorders and sensorineural deafness. Ocular

Background

collagen type XI alpha 1 chain(COL11A1) Homo sapiens This gene encodes one of the two alpha chains of type XI collagen, a minor fibrillar collagen. Type XI collagen is a heterotrimer but the third alpha chain is a post-translationally modified alpha 1 type II chain. Mutations in this gene are associated with type II Stickler syndrome and with Marshall syndrome. A single-nucleotide polymorphism in this gene is also associated with susceptibility to lumbar disc herniation. Multiple transcript variants have been identified for this gene. [provided by RefSeq, Nov 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using COL11A1 Monoclonal Antibody