



LI-cadherin Monoclonal Antibody

Catalog No	YP-mAb-17047
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CDH17
Immunogen	The antiserum was produced against synthesized peptide derived from human CDH17. AA range:341-390
Specificity	LI-cadherin Monoclonal Antibody detects endogenous levels of LI-cadherin protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	BILL-cadherin;CAD17_HUMAN;Cadherin 16;formerly;Cadherin 17;LI cadherin (liver-intestine);cadherin;liver-intestine;Cadherin-17;CDH16;Cdh17;FLJ26931; Formerly Cadherin 16;Formerly CDH16;HPT 1;HPT-1 cadherin;human intestinal peptide-associated transporter HPT-1;human peptide transporter 1;Intestinal peptide-associated transporter HPT 1;Intestinal peptide-associated transporter HPT-1;LI-cadherin;Liver Cadherin;liver intestine cadherin;Liver-intestine cadherin
Observed Band	99kD
Calculated Molecular Weight	99kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein .
Tissue Specificity	Expressed in the gastrointestinal tract and pancreatic duct. Not detected in kidney, lung, liver, brain, adrenal gland and skin.
Function	function:Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. LI-cadherin may have a role in the morphological organization of liver and intestine. Involved in intestinal peptide transport.,similarity:Contains 7 cadherin domains.,tissue specificity:Expressed in the gastrointestinal tract and pancreatic



duct. Not detected in kidney, lung, liver, brain, adrenal gland and skin.,

Background

This gene is a member of the cadherin superfamily, genes encoding calcium-dependent, membrane-associated glycoproteins. The encoded protein is cadherin-like, consisting of an extracellular region, containing 7 cadherin domains, and a transmembrane region but lacking the conserved cytoplasmic domain. The protein is a component of the gastrointestinal tract and pancreatic ducts, acting as an intestinal proton-dependent peptide transporter in the first step in oral absorption of many medically important peptide-based drugs. The protein may also play a role in the morphological organization of liver and intestine. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using LI-cadherin Monoclonal Antibody