



JAM-B Monoclonal Antibody

Catalog No	YP-mAb-17079
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	JAM2
Immunogen	Synthesized peptide derived from the Internal region of human JAM-B.
Specificity	JAM-B Monoclonal Antibody detects endogenous levels of JAM-B protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	JAMB;CD322;IBGC8;JAM-B;VEJAM;PRO245;VE-JAM;C21orf43;JAM2; Junctional adhesion molecule B;Junctional adhesion molecule 2;JAM-2;Vascular endothelial junction-associated molecule
Observed Band	33kD
Calculated Molecular Weight	33kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein . Cell junction . Cell junction, tight junction . Localized at tight junctions of both epithelial and endothelial cells (By similarity). Specifically localized within the somatodendritic compartment of neurons and excluded from the axon (By similarity). .
Tissue Specificity	Highly expressed in heart, placenta, lung, foreskin and lymph node (PubMed:10779521, PubMed:10945976). Prominently expressed on high endothelial venules and also present on the endothelia of other vessels (at protein level) (PubMed:10779521, PubMed:10945976). Also expressed in the brain in the caudate nuclei (PubMed:31851307).
Function	function:May play a role in the processes of lymphocyte homing to secondary lymphoid organs.,similarity:Belongs to the immunoglobulin superfamily., similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain., similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subcellular



location:Localized at tight junctions of both epithelial and endothelial cells.,
subunit:Interacts with JAM3.,tissue specificity:Highest expression in the heart,
placenta, lung, foreskin and lymph node. Prominently expressed on high
endothelial venules, also present on the endothelia of other vessels. Localized
to the intercellular boundaries of high endothelial cells.,

Background

This gene belongs to the immunoglobulin superfamily, and the junctional adhesion molecule (JAM) family. The protein encoded by this gene is a type I membrane protein that is localized at the tight junctions of both epithelial and endothelial cells. It acts as an adhesive ligand for interacting with a variety of immune cell types, and may play a role in lymphocyte homing to secondary lymphoid organs. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using JAM-B Monoclonal Antibody